

Nucleotide sequence of the EcoRI fragment from pLJ3 bearing two tandem lacUV5 promoters

Gregg Bogosian and James F.Kane

Animal Sciences Division, Monsanto Corporation, 700 Chesterfield Village Parkway, Chesterfield, MO 63198, USA

Submitted August 4, 1987

Accession no.Y00412

The plasmid pLJ3, a derivative of pMB9, contains a small EcoRI fragment bearing two tandem lacUV5 promoters (1). This EcoRI fragment was constructed in such a way that it consists of four segments: an EcoRI-AluI segment containing the first lacUV5 promoter, an AluI-HaeIII segment, a HaeIII-AluI segment, and an AluI-EcoRI segment containing the second lacUV5 promoter. All four segments were obtained from AluI fragments from λ plac5. The four AluI fragments were engineered via a complex series of steps to yield the EcoRI fragment of pLJ3 (1). pLJ3 has frequently been utilized as a source of the lacUV5 promoter for a variety of purposes (2-7). The nucleotide sequence of this EcoRI fragment was determined by the dideoxy method.

```

EcoRI                                     +1
GAATTCTCAC TCATTAGGCA CCCAGGCTT TACACTTTAT GCTTCCGGCT CGTATAATGT GTGGAATTGT
                                     PvuI/AluI       HaeIII
GAGCGGATAA CAATTTCACA CAGGAACAG CTGATTGCCC TTCACCGCCT GCCTCCGCTT GAGCCATCTG
                                     SphI         AluI
GATCGGCAGC GTTGTCTTCA TCAACCGGAA CGAGCATGCC GGAGAGCAGC TCACTCATTA GGCACCCAG
                                     +1
GCTTTAGACT TTATGCTTCC GGCTCGTATA ATGTGTGGAA TTGTGAGCGG ATAACAATTT CACACAGGAA
EcoRI
ACAGAATTC

```

Key restriction sites are shown above the DNA sequence, and +1 marks the startpoint of transcription for the two lacUV5 promoters. The 22 bp AluI-HaeIII segment is from the lacI gene (bp 1009 to 1030 of the Genbank sequence); the 67 bp HaeIII-AluI segment is presumably from a previously unsequenced portion of the *E. coli* chromosome carried on λ plac5. In addition to being a useful source of the lacUV5 promoter, we have found that this EcoRI fragment functions as an efficient transcription terminator. Others have reported transcription termination associated with the lac promoter (8-11).

References: (1) Johnsrud, L. (1978) Proc. Natl. Acad. Sci. USA 75: 5314, (2) Backman, K. and M. Ptashne (1978) Cell 13: 65, (3) Roberts, T. et al. (1979) Proc. Natl. Acad. Sci. USA 76: 760, (4) Guarente, L. et al. (1980) Cell 20: 543, (5) Lauer, G. et al. (1981) J. Mol. Appl. Gen. 1: 139, (6) Bogosian, G. and R. Somerville (1983) Mol. Gen. Genet. 191: 51, (7) Rasmussen, B. et al. (1985) J. Bacteriol. 164: 665, (8) Horowitz, H. and T. Platt (1982) J. Biol. Chem. 257: 11740, (9) Cone, K. et al. (1983) J. Biol. Chem. 258: 11296, (10) Deuschle, U. et al. (1986) Proc. Natl. Acad. Sci. USA 83: 4134, (11) Sellitti, M. et al. (1987) Proc. Natl. Acad. Sci. USA 84: 3199.